

# Graduate Algebra

Unless otherwise stated, all groups are finite, and all representations are of finite degree over  $\mathbb{C}$ .

♠ 2011/04/08

1, Let  $\rho : G \rightarrow GL_n(\mathbb{C})$  be a representation of a finite group  $G$ .

(i), Prove that  $\pi : g \mapsto \det \rho_g$  is one dimensional representation.

(ii), Prove that the quotient group  $G/\ker \pi$  is abelian.

2, Let  $S_n$  be the symmetric group on  $n$  letters. Prove that

(i),  $S_3 = \langle (12), (123) \rangle$ .

(ii), Denote by

$$\rho_{(12)} = \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix}, \quad \rho_{(123)} = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}.$$

Then  $\rho : S_3 \rightarrow GL_2(\mathbb{C})$  is a representation.

(iii),  $\rho$  is irreducible.

3, Let  $\rho : G \rightarrow GL(V)$  be a representation of a finite group  $G$ . Define the fixed subspace

$$V^G = \{v \in V \mid \rho_g v = v, \forall g \in G\}.$$

(i), Show that  $V^G$  is a  $G$ -invariant subspace of  $V$ .

(ii), Show that, for all  $v \in V$ , we have

$$\frac{1}{|G|} \sum_{h \in G} \rho_h v \in V^G.$$

(iii), Show that if  $v \in V^G$ , then

$$\frac{1}{|G|} \sum_{h \in G} \rho_h v = v.$$

♠ 2011/04/15

4, Let  $G$  be an abelian group. Then any irreducible representation of  $G$  has degree one.

5, Let  $\rho : G \rightarrow GL(V)$  be a representation of  $G$  with character  $\chi$  and let  $V^*$  be the dual space of  $V$ , i.e.,  $V^* = \text{Hom}(V, \mathbb{C})$ , the linear forms on  $V$ . For  $v \in V, v^* \in V^*$ , let  $\langle v, v^* \rangle = v^*(v)$ . Show that there exists a unique linear representation  $\rho^* : G \rightarrow GL(V^*)$ , such that

$$\langle \rho_g v, \rho_g^* v^* \rangle = \langle v, v^* \rangle, \quad \forall g \in G, v \in V, v^* \in V^*.$$

This is called the contragredient representation of  $\rho$  and its character is  $\bar{\chi}$ .

6, Let  $\rho : G \rightarrow GL(V)$  be an irreducible representation. Let

$$Z(G) = \{g \in G \mid gh = hg, \forall h \in G\}$$

be the center of  $G$ . Show that if  $g \in Z(G)$ , then  $\rho_g = \lambda I$  for some  $\lambda \in \mathbb{C}^\times$ .

♠ 2011/04/22

7, Let  $\chi$  be a nontrivial irreducible character of  $G$ . Show that

$$\sum_{g \in G} \chi(g) = 0.$$

8, Let  $\rho : S_n \rightarrow GL_n(\mathbb{C})$  be a representation given by defining  $\rho_\sigma(e_i) = e_{\sigma(i)}$  on the standard basis  $\{e_1, \dots, e_n\}$  for  $\mathbb{C}^n$ .

(i), Show that  $\chi_\rho(\sigma)$  is the number of fixed points of  $\sigma \in S_n$ , that is, the number of elements  $k \in \{1, \dots, n\}$  such that  $\sigma(k) = k$ .

(ii), Show that if  $n = 3$ , then  $\langle \chi_\rho, \chi_\rho \rangle = 2$  and hence  $\rho$  is not irreducible.

9, Let  $\rho : G \rightarrow GL_n(\mathbb{C})$  be a representation of  $G$  of dimension  $d$  and with character  $\chi$ . Show that  $\ker \rho = \{g \in G \mid \chi(g) = d\}$ . Show further that  $|\chi(g)| \leq d$  for all  $g \in G$ , with equality only if  $\rho(g) = \zeta I$ , a scalar multiple of the identity, for some root of unity  $\zeta$ .

Guanghua Ji · 纪广华  
School of Mathematics  
Shandong University  
Jinan, Shandong 250100  
[www.prime.sdu.edu.cn/ghji/guanghuaji.htm](http://www.prime.sdu.edu.cn/ghji/guanghuaji.htm)